

# Electroweak structure of light nuclei

Saori Pastore

Workshop on Progress in Ab Initio Techniques in Nuclear Physics  
@ TRIUMF, Vancouver BC, Canada - February 2016



\* in collaboration with \*

Rocco Schiavilla - JLab/ODU


Bob Wiringa, Steven Pieper, Maria Piarulli - ANL

Stefano Gandolfi, Joe Carlson - LANL

Luca Girlanda, Michele Viviani, Laura E. Marcucci, Alejandro Kievsky

- Salento U/INFN/Pisa U

## Electromagnetic probes to test predictive power of nuclear theories/models



\* Validate our theoretical understanding and control of nuclear **EM** structure and reactions is an essential prerequisite for studies on: \*

- ⇒ Weak induced reactions, *e.g.*,  $\nu$ -nucleus interactions (major progress by **A. Lovato**, **S. Gandolfi** *et al.*)
- ⇒ Larger nuclear systems


## The Basic Model: Nuclear Potentials

- ▶ The nucleus is a system made of  $A$  non-relativistic interacting nucleons, its energy is given by

$$H = T + V = \sum_{i=1}^A t_i + \sum_{i<j} v_{ij} + \sum_{i<j<k} V_{ijk} + \dots$$

where  $v_{ij}$  and  $V_{ijk}$  are 2- and 3-nucleon interaction operators

- ▶ Realistic  $v_{ij}$  and  $V_{ijk}$  interactions are based on EXPT data fitting and fitted parameters subsume underlying QCD
- ▶ Realistic potentials at large inter-particle distances are described in terms of one-pion-exchange, range  $\sim 1/m_\pi$ . Other mechanisms are, *e.g.*, two-pion exchange, range  $\sim 1/2m_\pi$ ;  $\Delta$ -excitations ...




- ▶ Potentials utilized in these sets of calculations to generate nuclear wave functions  $|\Psi_i\rangle$  solving  $H|\Psi_i\rangle = E_i|\Psi_i\rangle$  are:  
[AV18+UIX], [AV18+IL7], [NN(N3LO)+3N(N2LO)]

## The Basic Model: Nuclear Electromagnetic Currents - Impulse Approximation

- ▶ Current and charge operators describe the interaction of nuclei with external fields. They are expanded as a sum of 1-, 2-, ... nucleon operators:

$$\rho = \sum_{i=1}^A \rho_i + \sum_{i<j} \rho_{ij} + \dots, \quad \mathbf{j} = \sum_{i=1}^A \mathbf{j}_i + \sum_{i<j} \mathbf{j}_{ij} + \dots$$

- ▶ In Impulse Approximation **IA** nuclear EM currents are expressed in terms of those associated with individual protons and nucleons, *i.e.*,  $\rho_i$  and  $\mathbf{j}_i$




- ▶ IA picture is however incomplete; Historical evidence is the 10% underestimate of the  $np$  radiative capture ‘fixed’ by incorporating corrections from two-body meson-exchange EM currents - Riska&Brown 1972

## The Basic Model: Nuclear Electromagnetic Currents

- ▶ Current and charge operators describe the interaction of nuclei with external fields. They are expanded as a sum of 1-, 2-, ... nucleon operators:

$$\rho = \sum_{i=1}^A \rho_i + \sum_{i<j} \rho_{ij} + \dots, \quad \mathbf{j} = \sum_{i=1}^A \mathbf{j}_i + \sum_{i<j} \mathbf{j}_{ij} + \dots$$



- ▶ Longitudinal EM current operator  $\mathbf{j}$  linked to the nuclear Hamiltonian via continuity eq. ( $\mathbf{q}$  momentum carried by the external EM probe  $\gamma$ )

$$\mathbf{q} \cdot \mathbf{j} = [H, \rho] = [t_i + \mathbf{v}_{ij} + V_{ijk}, \rho]$$

- \* Meson-exchange currents **MEC** follow once meson-exchange mechanisms are implemented to describe nuclear forces - Villars&Miyazawa 40ies

These days we have:


- ▶ Highly sophisticated MEC projected out realistic potentials
- ▶ EM currents derived from  $\chi$ EFTs

## χEFT EM current up to $n = 1$ (or up to N3LO)

**LO** :  $j^{(-2)} \sim eQ^{-2}$



**NLO** :  $j^{(-1)} \sim eQ^{-1}$



**N<sup>2</sup>LO** :  $j^{(-0)} \sim eQ^0$



\* Two-body charge operators enter at N3LO and do not depend on LECs \*

▶ LO = IA

N2LO = IA(relativistic-correction)

▶ Strong contact LECs at N3LO fixed from fits to  $np$  phases shifts


PRC68, 041001 (2003)

▶ Unknown EM LECs enter the N3LO contact and tree-level currents


▶ No three-body EM currents at this order !!!

▶ NLO and N3LO loop-contributions lead to purely isovector operators

**N<sup>3</sup>LO** :  $j^{(1)} \sim eQ$




unknown LEC's →




PRC78(2008)064002, PRC80(2009)034004, PRC84(2011)024001

## $\chi$ EFT EM currents at N3LO: fixing the EM LECs



Five LECs:  $d^S$ ,  $d_1^V$ , and  $d_2^V$  could be determined by pion photo-production data on the nucleon



$d_2^V$  and  $d_1^V$  are known assuming  $\Delta$ -resonance saturation

Left with 3 LECs: Fixed in the  $A = 2 - 3$  nucleons' sector

▶ Isoscalar sector:

\*  $d^S$  and  $c^S$  from EXPT  $\mu_d$  and  $\mu_S(^3\text{H}/^3\text{He})$

▶ Isovector sector:

\*  $c^V$  from EXPT  $\mu_V(^3\text{H}/^3\text{He})$  m.m. ← our choice

Note that:



$\chi$ EFT operators have a power law behavior  $\rightarrow$  introduce a regulator to kill divergencies at large  $Q$ , e.g.,  $C_\Lambda = e^{-(Q/\Lambda)^n}$ , ...and also, pick  $n$  large enough so as to not generate spurious contributions

$$C_\Lambda \sim 1 - \left(\frac{Q}{\Lambda}\right)^n + \dots$$

Applications:  
EM form factors of nuclei with  $A = 2$  and  $3$




## Predictions with $\chi$ EFT EM Currents for the Deuteron Charge and Quadrupole f.f.'s



$\Lambda$ MeV	$\langle r_d \rangle$ (fm)	$\langle r_d \rangle$ EXP	$Q_d$ (fm <sup>2</sup> )	$Q_d$ (fm <sup>2</sup> ) EXP
500	1.976	1.9734(44)	0.285	0.2859(3)
600	1.968		0.282	


- ▶ Calculations include nucleonic f.f.'s taken from EXPT data
- ▶ Sensitivity to the cutoff used to regularize divergencies in the matrix elements is given by the bands' thickness

# Predictions with $\chi$ EFT EM Currents for the Deuteron Magnetic f.f.



PRC86(2012)047001 & PRC87(2013)014006

## Predictions with $\chi$ EFT EM Currents for ${}^3\text{He}$ and ${}^3\text{H}$ Magnetic f.f.'s



**LO/N3LO** with AV18+UIX – **LO/N3LO** with  $\chi$ -potentials NN(N3LO)+3N(N2LO)

- ▶  ${}^3\text{He}/{}^3\text{H}$  m.m.'s used to fix EM LECs;  $\sim 15\%$  correction from two-body currents
- ▶ Two-body corrections crucial to improve agreement with EXPT data

	${}^3\text{He} \langle r \rangle_{\text{EXP}} = 1.976 \pm 0.047 \text{ fm}$		${}^3\text{H} \langle r \rangle_{\text{EXP}} = 1.840 \pm 0.181 \text{ fm}$	
$\Lambda$	500	600	500	600
LO	2.098 (2.092)	2.090 (2.092)	1.924 (1.918)	1.914 (1.918)
N3LO	1.927 (1.915)	1.913 (1.924)	1.808 (1.792)	1.794 (1.797)

PRC87(2013)014006

## Benchmark calculations of $^3\text{He}$ Zemach Moments\*

**Quote:** Precise moments are useful observables for the comparison with theoretical calculations, ... in particular for light nuclei where very accurate *ab initio* calculations can be performed. I. Sick - [PRC90\(2014\)064002](#)

$$\langle r \rangle_{(2)} \propto - \int_0^\infty \frac{dq}{q^2} [G_E G_M - 1], \quad \langle r^3 \rangle_{(2)} \propto \int_0^\infty \frac{dq}{q^4} [G_E^2 - 1 + q^2 R^2 / 3]$$

	VMC(IA)	VMC(TOT)	GFMC(IA)	GFMC(TOT)	EXPT
$\langle r \rangle_{(2)}$	2.522	2.477	2.504	2.454	$2.528 \pm 0.016 \text{ fm}$
$\langle r^3 \rangle_{(2)}$	27.40	n.a.	29.30	n.a.	$28.15 \pm 0.70 \text{ fm}^3$
$\langle r_{\text{ch}}^2 \rangle^{1/2}$	1.967	n.a.	1.970	n.a.	$1.973 \pm 0.014 \text{ fm}$
$\langle r_{\text{m}}^2 \rangle^{1/2}$	2.000	1.962	2.019	1.942	$1.976 \pm 0.047 \text{ fm}$
$\langle r_{\text{ch}}^4 \rangle$	19.8	n.a.	30.0	n.a.	$32.9 \pm 1.60 \text{ fm}^4$
$\langle \mu \rangle$	-1.775	-2.134	-1.767	-2.129	$-2.127 \mu_N$

\* in collaboration with

Nir Nievo, Chen Ji, Sonia Bacca, Maria Piarulli and Bob Wiringa

Preliminary!!!

## Calculations with EM Currents from $\chi$ EFT with $\pi$ 's and N's

- ▶ Park, Min, and Rho *et al.* (1996)

applications to:

magnetic moments and M1 properties of  $A=2-3$  systems, and radiative captures in  $A=2-4$  systems by Song, Lazauskas, Park *et al.* (2009-2011) within the hybrid approach

.....

\* Based on EM  $\chi$ EFT currents from [NPA596\(1996\)515](#)

- ▶ Meissner and Walzl (2001);

Kölling, Epelbaum, Krebs, and Meissner (2009-2011)

applications to:

$d$  and  $^3\text{He}$  photodisintegration by Rozpedzik *et al.* (2011);  $e$ -scattering (2014);

$d$  magnetic f.f. by Kölling, Epelbaum, Phillips (2012);

radiative  $N-d$  capture by Skibinski *et al.* (2014)

.....

\* Based on EM  $\chi$ EFT currents from [PRC80\(2009\)045502](#) & [PRC84\(2011\)054008](#) and consistent  $\chi$ EFT potentials from UT method

- ▶ Phillips (2003-2007)


applications to [deuteron static properties and f.f.'s](#)

.....

Moving on to larger nuclear systems:  
magnetic moments and transitions in  $A \leq 10$  nuclei

# Magnetic Moments in $A \leq 10$ Nuclei



## Predictions for $A > 3$ nuclei



- ▶  $\mu(\text{IA}) = \mu_N \sum_i [(L_i + g_p S_i)(1 + \tau_{i,z})/2 + g_n S_i(1 - \tau_{i,z})/2]$
- ▶ GFMC calculations based on  $H = T + \text{AV18} + \text{IL7}$

# Magnetic Moments in $A \leq 10$ Nuclei - bis

## Predictions for $A > 3$ nuclei



- ▶  $\mu_N(\text{IA}) = \sum_i [(L_i + g_p S_i)(1 + \tau_{i,z})/2 + g_n S_i(1 - \tau_{i,z})/2]$
- ▶  ${}^9\text{C}$  ( ${}^9\text{Li}$ ) dominant spatial symmetry [s.s.] = [432] =  $[\alpha, {}^3\text{He}({}^3\text{H}), pp(nn)] \rightarrow$  Large MEC
- ▶  ${}^9\text{Be}$  ( ${}^9\text{B}$ ) dominant spatial symmetry [s.s.] = [441] =  $[\alpha, \alpha, n(p)]$

PRC87(2013)035503



## EM Transitions in $A \leq 9$ Nuclei

- ▶ Two-body EM currents bring the theory in a better agreement with the EXP
- ▶ Significant correction in  $A = 9$ ,  $T = 3/2$  systems. Up to  $\sim 40\%$  correction found in  ${}^9\text{C}$  m.m.
- ▶ Major correction ( $\sim 60 - 70\%$  of total MEC) is due to the one-pion-exchange currents at NLO – purely isovector


One M1 prediction:  ${}^9\text{Li}(1/2 \rightarrow 3/2)^*$

$$\Gamma(\text{IA}) = 0.59(2) \text{ eV}$$

$$\Gamma(\text{TOT}) = 0.79(3) \text{ eV}$$


+ a number of B(E2)s in IA

\*Ricard-McCutchan *et al.* TRIUMF proposal 2014 - ongoing data analysis



# $^8\text{Be}$ Energy Spectrum


- ▶  $2^+$  and  $4^+$  broad states at  $\sim 3$  MeV and  $\sim 11$  MeV
- ▶ isospin-mixed states at  $\sim 16$  MeV,  $\sim 17$  MeV,  $\sim 19$  MeV
- ▶ **M1** transitions
- ▶ **E2** transitions
- ▶ **E2 + M1** transitions



$J^\pi; T$	GFMC	Iso-mixed	Experiment
$0^+$	-56.3(1)		-56.50
$2^+$	+ 3.2(2)	+ 3.03(1)	+ 3.03(1)
$4^+$	+11.2(3)		+11.35(15)
$2^+; 0$	+16.8(2)	+16.746(3)	+16.626(3)
$2^+; 1$	+16.8(2)	+16.802(3)	+16.922(3)
$1^+; 1$	+17.5(2)	+17.67	+17.640(1)
$1^+; 0$	+18.0(2)	+18.12	+18.150(4)
$3^+; 1$	+19.4(2)	+19.10	+19.07(3)
$3^+; 0$	+19.9(2)	+19.21	+19.235(10)

PRL111(2013)062502 & PRC90(2014)024321


## Two-body M1 transitions densities



$(J_i, T_i) \rightarrow (J_f, T_f)$	IA	NLO-OPE	N2LO-RC	N3LO-TPE	N3LO-CT	N3LO- $\Delta$	MEC
$(1^+; 1) \rightarrow (2_2^+; 0)$	2.461 (13)	0.457 (3)	-0.058 (1)	0.095 (2)	-0.035 (3)	0.161 (21)	0.620 (5)

PRC90(2014)024321


## M1 Transition Widths / EXPT



- ▶ Predictions for [s.s.]-conserving transitions are in fair agreement with EXPT
- ▶ The theoretical description for this system is unsatisfactory, however, MEC provide a  $\sim 20 - 30\%$  correction to the calculated matrix elements improving the agreement with EXPT data

Beta-decay rates for  $A \leq 10$  nuclei

## Theory vs Experiment: Quenching



$$3 \leq A \leq 18$$

Fig. from Chou *et al.* PRC47(1993)163

perfect agreement

theory > experiment

temporary fix:

$$g_A^{\text{eff}} \simeq 0.70 g_A$$

Quenching origin: *i*) better w.f.'s and/or *ii*) many body currents are required

$\beta \pm - (J_i^\pi, T_i) \rightarrow (J_f^\pi, T_f)$	simple w.f.'s	IA	IA+MEC	Experiment
${}^3\text{H}(1/2^+, 1/2) \rightarrow {}^3\text{He}(1/2^+, 1/2)$	2.449	2.2765(1)		2.357(10)*
${}^6\text{He}(0^+, 1) \rightarrow {}^6\text{Li}(1^+, 0)$	2.449	2.150	2.187	2.182*
${}^7\text{Be}(3/2^-, 1/2) \rightarrow {}^7\text{Li}(3/2^-, 1/2)$	2.582	2.292	2.395	2.290*
${}^{10}\text{C}(0^+, 1) \rightarrow {}^{10}\text{B}(1^+, 0)$	2.449	2.024	2.076	1.862*

Preliminary!!!

• in collaboration with **Bob Wiringa, Stefano Gandolfi and Rocco Schiavilla**

\* data from TUNL compilations

\* data from Suzuki *et al.* PRC67(2003)044302

## Summary

The microscopic description of nuclei successfully reproduces EXPT data provided that many-body effects in nuclear interactions and EM currents are accounted for.

J.Phys.G41(2014)123002 - S.Bacca&S.P.

- ▶ Two-body EM currents from  $\chi$ EFT tested in  $A \leq 10$  nuclei
- ▶ Two-body corrections can be sizable and improve on theory/EXPT agreement
- ▶ EM structure of  $A = 2-3$  nuclei well reproduced with chiral charge and current operators for  $q \lesssim 3m_\pi$
- ▶  $\sim 40\%$  two-body correction found in  ${}^9\text{C}$ 's m.m.
- ▶  $\sim 20-30\%$  corrections found in M1 transitions in low-lying states of  ${}^8\text{Be}$
- ▶  $\sim 10\%$  contributions from correlations in beta-decay m.e.'s in  $A \leq 10$  nuclei

## Outlook

The microscopic description of nuclei successfully reproduces EXPT data provided that many-body effects in nuclear interactions and EM currents are accounted for.


J.Phys.G41(2014)123002 - S.Bacca&S.P.

- \* EM structure and dynamics of light nuclei
  - ▶ Charge and magnetic form factors of  $A \leq 10$  systems
  - ▶ M1/E2 transitions in light nuclei
  - ▶ Radiative captures, photonuclear reactions . . .
  - ▶ Fully consistent  $\chi$ EFT calculations with ‘MEC’ for  $A > 4$
  - ▶ Role of  $\Delta$ -resonances in ‘MEC’ (EM current consistent with the chiral ‘ $\Delta$ -full’ NN potential developed by M. Piarulli et al. PRC91(2015)024003)
- \* Electroweak structure and dynamics of light nuclei
  - ▶  $\nu$ -nucleus scattering J. Carlson, S. Gandolfi, B. Wiringa, R. Schaivilla
  - ▶ Test axial currents (chiral and conventional) in light nuclei (A. Baroni et al. PRC93(2016)015501)
  - ▶ Many-body effects in  $\nu$ - $d$  pion-production at threshold (in preparation)



## EXTRA SLIDES

## NN Potential at NLO (or $Q^{n=2}$ )




- ▶ Contact potential at LO (or  $Q^{n=0}$ ) depends on 2 LECs
- ▶ Contact potential at NLO (or  $Q^{n=2}$ ) depends on 7 additional LECs


### NN potentials with $\pi$ 's and $N$ 's

- \* van Kolck *et al.* (1994–96)
- \* Kaiser, Weise *et al.* (1997–98)
- \* Epelbaum, Glöckle, Meissner (1998–2015)
- \* Entem and Machleidt (2002–2015) ←
- \* ...

## $\chi$ EFT EM currents at N3LO: fixing the EM LECs



Five LECs:  $d^S$ ,  $d_1^V$ , and  $d_2^V$  could be determined by pion photo-production data on the nucleon



Isovector  $d_2^V$  and  $d_1^V$  are known assuming  $\Delta$ -resonance saturation

Left with 3 LECs: Fixed in the  $A = 2 - 3$  nucleons' sector

▶ Isoscalar sector:

\*  $d^S$  and  $c^S$  from EXPT  $\mu_d$  and  $\mu_S(^3\text{H}/^3\text{He})$

▶ Isovector sector:

\* model I =  $c^V$  from EXPT  $npd\gamma$  xsec.

or

\* model II =  $c^V$  from EXPT  $\mu_V(^3\text{H}/^3\text{He})$  m.m. ← our choice


Note that:

$\chi$ EFT operators have a power law behavior  $\rightarrow$  introduce a regulator to kill divergencies at large  $Q$ , e.g.,  $C_\Lambda = e^{-(Q/\Lambda)^n}$ , ...and also, pick  $n$  large enough so as to not generate spurious contributions

$$C_\Lambda \sim 1 - \left(\frac{Q}{\Lambda}\right)^n + \dots$$


## Predictions with $\chi$ EFT EM currents for $A = 2-3$ systems

$np$  capture xsec. (using model II) /  $\mu_V$  of  $A = 3$  nuclei (using model I)  
bands represent nuclear model dependence (N3LO/N2LO – AV18/UIX)



- ▶  $npd\gamma$  xsec. and  $\mu_V(^3\text{H}/^3\text{He})$  m.m. are within 1% and 3% of EXPT
- ▶ Two-body currents important to reach agreement with exp data
- ▶ Negligible dependence on the cutoff entering the regulator  $\exp(-(k/\Lambda)^4)$


## Deuteron wave functions



from Entem&Machleidt 2011 Review

- ▶ Entem&Machleidt N3LO
- ▶ Epelbaum *et al.* 2005
- ▶ black lines = conventional potentials, *i.e.* AV18, CD-Bonn, Nijm-I

## $^3\text{He}$ and $^3\text{H}$ charge f.f.'s



- ▶ Excellent agreement up to  $q \simeq 2 \text{ fm}^{-1}$
- ▶ N3LO and N4LO comparable

	$^3\text{He} \langle r \rangle_{\text{EXP}} = 1.959 \pm 0.030 \text{ fm}$		$^3\text{H} \langle r \rangle_{\text{EXP}} = 1.755 \pm 0.086$	
$\Lambda$	500	600	500	600
LO	1.966 (1.950)	1.958 (1.950)	1.762 (1.743)	1.750 (1.743)
N4LO	1.966 (1.950)	1.958 (1.950)	1.762 (1.743)	1.750 (1.743)


## E2 transitions in ${}^8\text{Be}$

- ▶  $2^+$  and  $4^+$  broad rotational states at  $\sim 3$  MeV and  $\sim 11$  MeV
- ▶  $4^+ \rightarrow 2^+$  transition recently measured at BARC\*, Mumbai
- ▶ Computational challenge:  $2^+$  and  $4^+$  states tend to break up into two  $\alpha$  as  $\tau$  increases
- ▶ Results obtained by linear fitting the GFMC points and extrapolating at  $\tau = 0.1$  MeV where stability is observed in the g.s. energy propagation

$J^\pi; T$	$E$ [MeV]	$B(E2)$ [ $e^2 \text{fm}^4$ ]
$0^+$	-56.3(1)	
$2^+$	+3.2(2)	20.0 (8)- [ $2^+ \rightarrow 0^+$ ]
$4^+$	+11.2(3)	27.2(15)- [ $4^+ \rightarrow 2^+$ ]*


\*Bhabha Atomic Research Centre

\*EXPT  $B(E2) = 21 \pm 2.3 e^2 \text{fm}^4$



## E2 transition widths / EXPT

- ▶ We attempt to evaluate a number of E2 transitions (predictions not shown in the figure)
- ▶ Complications are due to large cancellations among large m.e.'s  $\rightarrow$  E2s very sensitive to small components
- ▶ One more complication: make sure that the first and second  $(J^\pi, T) = (2^+, 0)$  states are orthogonal



- \* We orthogonalize the second  $(J^\pi, T) = (2^+, 0)$  via

$$|\Psi_{2^+}^+(\text{ortho})\rangle_G = |\Psi_{2^+}^+\rangle_G - \langle \Psi_{2^+}^+ | \Psi_{2^+}^+ \rangle_V |\Psi_{2^+}^+\rangle_G$$



## Anomalous magnetic moment of ${}^9\text{C}$

### Mirror nuclei spin expectation value

- ▶ Charge Symmetry Conserving (CSC) picture ( $p \leftrightarrow n$ )  $\diamond$

$$\langle \sigma_z \rangle = \frac{\mu(T_z = +T) + \mu(T_z = -T) - J}{(g_s^p + g_s^n - 1)/2} = \frac{2\mu(\text{IS}) - J}{0.3796}$$

- ▶ For  $A = 9$ ,  $T = 3/2$  mirror nuclei:  ${}^9\text{C}$  and  ${}^9\text{Li}$   
 EXP  $\langle \sigma_z \rangle = 1.44$  while THEORY  $\langle \sigma_z \rangle \sim 1$  (assuming CSC)  
 possible cause: Charge Symmetry Breaking (CSB)
- ▶ Three different predictions for  $\langle \sigma_z \rangle$  with CSC w.f.'s (\*) and CSB w.f.'s

$\langle \sigma_z \rangle$	Symmetry	IA	TOT	EXP
CSB	${}^9\text{Li}(\frac{3}{2}^-, \frac{3}{2}^-), {}^9\text{C}(\frac{3}{2}^-, \frac{3}{2}^-)$	1.05(1)	1.31(11)	1.44
CSC	${}^9\text{Li}(\frac{3}{2}^-, \frac{3}{2}^-), {}^9\text{C}(\frac{3}{2}^-, \frac{3}{2}^-)^*$	0.95 (11)	1.00 (11)	
CSC	${}^9\text{Li}(\frac{3}{2}^-, \frac{3}{2}^-)^*, {}^9\text{C}(\frac{3}{2}^-, \frac{3}{2}^-)$	1.00 (1)	1.05 (1)	

- ▶ **Need both CSB in the w.f.'s and MEC!**

$\diamond$  Utsuno – PRC70, 011303(R) (2004)

## Currents from nuclear interactions \*- Marcucci *et al.* PRC72, 014001 (2005)

- ▶ Current operator  $\mathbf{j}$  constructed so as to satisfy the continuity equation with a realistic Hamiltonian
- ▶ Short- and intermediate-behavior of the EM operators inferred from the nuclear two- and three-body potentials

$$\mathbf{j} = \mathbf{j}^{(1)} + \mathbf{j}^{(2)}(v) + \mathbf{j}^{(3)}(V)$$


The diagram illustrates two Feynman diagrams for nuclear current operators. The first diagram shows a nucleon (N) emitting a pion ( $\pi$ ) and then interacting with another nucleon (N) via a  $\Delta$  resonance and a wavy line representing a pion with momentum  $q$ . The second diagram shows a nucleon (N) interacting with another nucleon (N) via a pion ( $\pi$ ) and a  $\rho$  meson with momentum  $\omega$ . A bracket above the diagrams is labeled "transverse".

\* also referred to as Standard Nuclear Physics Approach (SNPA) currents

- ▶ Long range part of  $\mathbf{j}(v)$  corresponds to OPE seagull and pion-in-flight EM currents

## Currents from nuclear interactions - Marcucci *et al.* PRC72, 014001 (2005)

Satisfactory description of a variety of nuclear EM properties [see Marcucci *et al.* (2005) and (2008)]



- ▶ Isoscalar magnetic moments are a few % off (10% in  $A=7$  nuclei)

## Magnetic moments in $A \leq 9$ nuclei: SNPA vs $\chi$ EFT

	A	s.s.	IA	TOT SNPA	TOT $\chi$ EFT*	EXP
IS	7	[43]	0.902 (3)	0.833 (12)	0.906 (7)	0.929
IV		[43]	-3.944 (5)	-4.587 (18)	-4.670 (9)	-4.654
IS	8	[431]	1.289 (8)	1.160 (15)	1.299 (9)	1.344
IV		[431]	0.182 (8)	-0.129 (15)	-0.139 (9)	-0.310
IS	9	[432]	0.994 (15)	0.922 (32)	1.038 (21)	1.024
IV		[432]	-1.095 (10)	-1.371 (21)	-1.532 (15)	-1.610

### Preliminary results

Overall improvement of isoscalar (IS) component of the magnetic moment

$$\mu = \mu_S + \tau_z \mu_V$$

## NUCLEAR HAMILTONIAN

$$H = \sum_i K_i + \sum_{i < j} v_{ij} + \sum_{i < j < k} V_{ijk}$$

$K_i$ : Non-relativistic kinetic energy,  $m_n$ - $m_p$  effects included


Argonne v<sub>18</sub>:  $v_{ij} = v_{ij}^T + v_{ij}^\pi + v_{ij}^I + v_{ij}^S = \sum v_p(r_{ij}) O_{ij}^p$

- 18 spin, tensor, spin-orbit, isospin, etc., operators
- full EM and strong CD and CSB terms included
- predominantly local operator structure
- fits Nijmegen PWA93 data with  $\chi^2/\text{d.o.f.}=1.1$

Wiringa, Stoks, & Schiavilla, PRC **51**, (1995)


Urbana & Illinois:  $V_{ijk} = V_{ijk}^{2\pi} + V_{ijk}^{3\pi} + V_{ijk}^R$

- Urbana has standard  $2\pi$   $P$ -wave + short-range repulsion for matter saturation
- Illinois adds  $2\pi$   $S$ -wave +  $3\pi$  rings to provide extra  $T=3/2$  interaction
- Illinois-7 has four parameters fit to 23 levels in  $A \leq 10$  nuclei



Pieper, Pandharipande, Wiringa, & Carlson, PRC **64**, 014001 (2001)

Pieper, AIP CP **1011**, 143 (2008)




### THREE-NUCLEON POTENTIALS

$$\text{Urbana } V_{ijk} = V_{ijk}^{2\pi P} + V_{ijk}^R$$



Carlson, Pandharipande, & Wiringa, NP **A401**, 59 (1983)

$$\text{Illinois } V_{ijk} = V_{ijk}^{2\pi P} + V_{ijk}^{2\pi S} + V_{ijk}^{3\pi\Delta R} + V_{ijk}^R$$



Pieper, Pandharipande, Wiringa, & Carlson, PRC **64**, 014001 (2001)

**Illinois-7** has 4 strength parameters fit to 23 energy levels in  $A \leq 10$  nuclei.

In light nuclei we find (thanks to large cancellation between  $\langle K \rangle$  &  $\langle v_{ij} \rangle$ ):


$$\langle V_{ijk} \rangle \sim (0.02 \text{ to } 0.07) \quad \langle v_{ij} \rangle \sim (0.15 \text{ to } 0.5) \quad \langle H \rangle$$

We expect  $\langle \mathcal{V}_{ijkl} \rangle \sim 0.05$   $\langle V_{ijk} \rangle \sim (0.01 \text{ to } 0.03)$   $\langle H \rangle \sim 1 \text{ MeV}$  in  $^{12}\text{C}$ .

## Transition amplitude in time-ordered perturbation theory

$$\begin{aligned}
 T_{fi} = \langle f | T | i \rangle &= \langle f | H_1 \sum_{n=1}^{\infty} \left( \frac{1}{E_i - H_0 + i\eta} H_1 \right)^{n-1} | i \rangle \\
 &= \langle f | H_1 | i \rangle + \sum_{|I\rangle} \langle f | H_1 | I \rangle \frac{1}{E_i - E_I} \langle I | H_1 | i \rangle + \dots
 \end{aligned}$$

- ▶ A contribution with N interaction vertices and L loops scales as

$$\underbrace{e \left( \prod_{i=1}^N Q^{\alpha_i - \beta_i/2} \right)}_{H_1 \text{ scaling}} \times \underbrace{Q^{-(N-N_K-1)} Q^{-2N_K}}_{\text{denominators}} \times \underbrace{Q^{3L}}_{\text{loop integration}}$$


$\alpha_i$  = number of derivatives in  $H_1$  and  $\beta_i$  = number of  $\pi$ 's at each vertex

$N_K$  = number of pure nucleonic intermediate states


- ▶ Due to the chiral expansion, the transition amplitude  $T_{fi}$  can be expanded as

$$T_{fi} = T^{\text{LO}} + T^{\text{NLO}} + T^{\text{N}^2\text{LO}} + \dots \quad \text{and} \quad T^{\text{N}^n\text{LO}} \sim (Q/\Lambda_\chi)^n T^{\text{LO}}$$

## Power counting

- ▶  $N_K$  energy denominators scale as  $Q^{-2}$

$$\frac{1}{E_i - E_I} |I\rangle = \frac{1}{E_i - E_N} |I\rangle \sim Q^{-2} |I\rangle$$



- ▶  $(N - N_K - 1)$  energy denominators scale  $Q^{-1}$  in the static limit; they can be further expanded in powers of  $(E_i - E_N)/\omega_\pi \sim Q$

$$\frac{1}{E_i - E_I} |I\rangle = \frac{1}{E_i - E_N - \omega_\pi} |I\rangle \sim - \left[ \underbrace{\frac{1}{\omega_\pi}}_{Q^{-1}} + \underbrace{\frac{E_i - E_N}{\omega_\pi^2}}_{Q^0} + \underbrace{\frac{(E_i - E_N)^2}{\omega_\pi^3}}_{Q^1} + \dots \right] |I\rangle$$

- ▶ Terms accounted into the Lippmann-Schwinger Eq. are subtracted from the reducible amplitude
- ▶ EM operators depend on the off-the-energy shell prescription adopted for the non-static OPE and TPE potentials
- ▶ Ultimately, the EM operators are unitarily equivalent: Description of physical systems is not affected by this ambiguity



## Magnetic moment at N<sup>3</sup>LO

- ▶ Magnetic moment operator due to two-body current density  $\mathbf{J}(\mathbf{x})$

$$\boldsymbol{\mu}(\mathbf{R}, \mathbf{r}) = \frac{1}{2} \left[ \mathbf{R} \times \int d\mathbf{x} \mathbf{J}(\mathbf{x}) + \int d\mathbf{x} (\mathbf{x} - \mathbf{R}) \times \mathbf{J}(\mathbf{x}) \right]$$

- ▶ Sachs' and translationally invariant magnetic moments

$$\begin{aligned} \boldsymbol{\mu}_{\text{Sachs}}(\mathbf{R}, \mathbf{r}) &= -i \frac{\mathbf{R}}{2} \times \int d\mathbf{x} \mathbf{x} [\rho(\mathbf{x}), v_{12}] \\ \boldsymbol{\mu}_{\text{T}}(\mathbf{r}) &= -\frac{i}{2} \int_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} \nabla_q \times \mathbf{j}(\mathbf{q}, \mathbf{k}) \Big|_{\mathbf{q}=0} \end{aligned}$$

## 2009 EM current vs 2011 EM currents p. 1/2



g)

i)

- ▶ Non-static corrections entering single-nucleon operators accounted into the derivation of current i)

$$\begin{aligned}
 i) \text{ OLD} &= i \frac{e g_A^2}{F_\pi^2} \tau_{1,z} \int \frac{\mathbf{q}_1 - \mathbf{q}_2}{\omega_1^3 \omega_2^3} \frac{\omega_1^2 + \omega_1 \omega_2 + \omega_2^2}{\omega_1 + \omega_2} \left[ C_S \boldsymbol{\sigma}_1 \cdot (\mathbf{q}_1 \times \mathbf{q}_2) \right. \\
 &\quad \left. - C_T \boldsymbol{\sigma}_2 \cdot (\mathbf{q}_1 \times \mathbf{q}_2) \right] + 1 \rightleftharpoons 2 \\
 i) \text{ NEW} &= 2i \frac{e g_A^2 C_T}{F_\pi^2} \tau_{1,z} \int_{\mathbf{q}_1, \mathbf{q}_2} \frac{\omega_1^2 + \omega_1 \omega_2 + \omega_2^2}{\omega_1^3 \omega_2^3 (\omega_1 + \omega_2)} (\mathbf{q}_1 - \mathbf{q}_2) \boldsymbol{\sigma}_2 \cdot \mathbf{q}_2 \times \mathbf{q}_1 + 1 \rightleftharpoons 2
 \end{aligned}$$

- ▶ i) NEW in agreement with Kölling 2009/2011\* but for a factor of 2, which has no impact because  $(i + g) = 0$

\* PRC80, 045502 (2009)/ PRC 84, 054008 (2011)

## 2009 EM current vs 2011 EM currents p. 2/2



- ▶ A different derivation in Kölling 2009/2011\* leads to an additional term  $\sim (\boldsymbol{\sigma}_i \times \mathbf{q}) \times \mathbf{q}$  in the N3LO current at tree level, which however does not contribute to the magnetic moment
- ▶ The N3LO contact current of Pastore 2009 is in agreement with that of Kölling 2011 after Fierz-reordering, apart from differences in the term  $\propto C_5$ :

$$\mathbf{j}_{\text{ct}}^{\text{N3LO}} = -\frac{iC_5}{4} (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \times (e_1 \mathbf{k}_1 + e_2 \mathbf{k}_2)$$

\* PRC80, 045502 (2009)/ PRC 84, 054008 (2011)

## Magnetic moment (m.m.) operator

- ▶ comparison with Kölling *et al.*:
  - i) LO, NLO, N2LO, N3LO TPE, N3LO CT, N3LO TREE m.m.'s agree, but for the N3LO CT term  $\propto C_5$
  - ii) currents associated with one loop corrections to the OPE are missing in these calculations of m.m.'s; renormalization of OPE currents has been carried out in Kölling 2011\*
  
- ▶ comparison with Park *et al.*\*\*:
  - i) Sachs' m.m. is missing (no problem in two-body systems),
  - ii) TPE box contribution at N3LO generates an extra term  $\propto (\boldsymbol{\tau}_i \times \boldsymbol{\tau}_j)_z$
  
- \* Ultimately, in actual calculations these differences are presumably mitigated by fitting LECs to experimental data

\* loop corrections to OPE: terms  $\propto L(k)$  in Eq. (4.28) of Kölling 2011;


\*\* NP **A596**, 515, (1996)

## EM current up to $n = 1$ (or up to N3LO) - bis

**LO** :  $j^{(-2)} \sim eQ^{-2}$



**NLO** :  $j^{(-1)} \sim eQ^{-1}$




**N<sup>2</sup>LO** :  $j^{(-0)} \sim eQ^0$




- ▶  $n = -2, -1, 0,$  and 1-(loops only): depend on known LECs namely  $g_A, F_\pi,$  and proton and neutron  $\mu$
- ▶  $n = 0$ :  $(Q/m_N)^2$  relativistic correction to  $\mathbf{j}^{(-2)}$
- ▶ unknown LECs enter the  $n = 1$  contact and tree-level currents (the latter originates from a  $\gamma\pi N$  vertex of order  $eQ^2$ )

- ▶ divergencies associated with loop integrals are reabsorbed by renormalization of contact terms
- ▶ loops contributions lead to purely isovector operators
- ▶  $\mathbf{j}^{(n \leq 1)}$  satisfies the CCR with  $\chi$ EFT two-nucleon potential  $v^{(n \leq 2)}$

**N<sup>3</sup>LO**:  $j^{(1)} \sim eQ$



unknown LEC's →



$$v^{\text{ME}} = f_{\text{PS}} \left( \text{Diagram 1} \right) + \left( \text{Diagram 2} \right)$$

- ▶ Exploiting the meson exchange (ME) mechanism, one assumes that the static part  $v_0$  of  $v$  is due to pseudoscalar (PS) and vector (V) exchanges
- ▶  $v^{\text{ME}}$  is expressed in terms of 'effective propagators'  $v_{PS}$ ,  $v_V$ ,  $v_{VS}$ , fixed such to reproduce  $v_0$ , for example


$$v_{PS} = [v^{\sigma\tau}(k) - 2v^{t\tau}(k)]/3$$

with  $v^{\sigma\tau}$  and  $v^{t\tau}$  components of  $v_0$

- ▶ The current operator is obtained by taking the non relativistic reduction of the ME Feynman amplitudes and replacing the bare propagators with the 'effective' ones

$$j^{(2)}(v_0) = \left( \text{Diagram 1} \right) + \left( \text{Diagram 2} \right) + \left( \text{Diagram 3} \right)$$

## OPEP beyond the static limit



On-the-energy-shell, non-static OPEP at N2LO ( $Q^2$ ) can be equivalently written as

$$\begin{aligned}
 \mathbf{v}_{\pi}^{(2)}(\mathbf{v} = 0) &= \mathbf{v}_{\pi}^{(0)}(\mathbf{k}) \frac{(E'_1 - E_1)^2 + (E'_2 - E_2)^2}{2\omega_k^2} \\
 \mathbf{v}_{\pi}^{(2)}(\mathbf{v} = 1) &= -\mathbf{v}_{\pi}^{(0)}(\mathbf{k}) \frac{(E'_1 - E_1)(E'_2 - E_2)}{\omega_k^2} \\
 \mathbf{v}_{\pi}^{(0)}(\mathbf{k}) &= -\frac{g_A^2}{F_{\pi}^2} \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 \frac{\boldsymbol{\sigma}_1 \cdot \mathbf{k} \boldsymbol{\sigma}_2 \cdot \mathbf{k}}{\omega_k^2}
 \end{aligned}$$

$\mathbf{v}_{\pi}^{(2)}(\mathbf{v})$  corrections are different off-the-energy-shell ( $E_1 + E_2 \neq E'_1 + E'_2$ )

- ▶ TPE contributions are affected by the choice made for the parameter  $\mathbf{v}$

## From amplitudes to potentials

The two-nucleon potential  $v = v^{(0)} + v^{(1)} + v^{(2)} + \dots$  (with  $v^{(n)} \sim Q^n$ ) is iterated into the Lippmann-Schwinger (LS) equation *i.e.*


$$v + v G_0 v + v G_0 v G_0 v + \dots, \quad G_0 = 1/(E_i - E_I + i\eta)$$

$v^{(n)}$  is obtained subtracting from the transition amplitude  $T_{\text{fi}}^{(n)}$  terms already accounted for into the LS equation

$$\begin{aligned} v^{(0)} &= T^{(0)}, \\ v^{(1)} &= T^{(1)} - \left[ v^{(0)} G_0 v^{(0)} \right], \\ v^{(2)} &= T^{(2)} - \left[ v^{(0)} G_0 v^{(0)} G_0 v^{(0)} \right] - \left[ v^{(1)} G_0 v^{(0)} + v^{(0)} G_0 v^{(1)} \right], \\ v^{(3)}(\mathbf{v}) &= T^{(3)} - \left[ v^{(0)} G_0 v^{(0)} G_0 v^{(0)} G_0 v^{(0)} \right] - \left[ v^{(1)} G_0 v^{(0)} G_0 v^{(0)} + \text{permutations} \right] \\ &\quad - \underbrace{\left[ v^{(1)} G_0 v^{(1)} \right] - \left[ v^{(2)}(\mathbf{v}) G_0 v^{(0)} + v^{(0)} G_0 v^{(2)}(\mathbf{v}) \right]}_{\text{LS terms}} \end{aligned}$$



# From amplitudes to potentials: an example with OPE and TPE only



► To each  $v_\pi^{(2)}(\mathbf{v})$  corresponds a  $v_{2\pi}^{(3)}(\mathbf{v})$

## Unitary equivalence of $v_{\pi}^{(2)}(\mathbf{v})$ and $v_{2\pi}^{(3)}(\mathbf{v})$

- ▶ Different off-the-energy-shell parameterizations lead to unitarily equivalent two-nucleon Hamiltonians

$$H(\mathbf{v}) = t^{(-1)} + v_{\pi}^{(0)} + v_{2\pi}^{(2)} + v_{\pi}^{(2)}(\mathbf{v}) + v_{2\pi}^{(3)}(\mathbf{v})$$

$t^{(-1)}$  is the kinetic energy,  $v_{\pi}^{(0)}$  and  $v_{2\pi}^{(2)}$  are the static OPEP and TPEP

- ▶ The Hamiltonians are related to each other via

$$H(\mathbf{v}) = e^{-iU(\mathbf{v})} H(\mathbf{v} = 0) e^{+iU(\mathbf{v})}, \quad iU(\mathbf{v}) \simeq iU^{(0)}(\mathbf{v}) + iU^{(1)}(\mathbf{v})$$

from which it follows

$$H(\mathbf{v}) = H(\mathbf{v} = 0) + \left[ t^{(-1)} + v_{\pi}^{(0)}, iU^{(0)}(\mathbf{v}) \right] + \left[ t^{(-1)}, iU^{(1)}(\mathbf{v}) \right]$$

- ▶ Predictions for physical observables are unaffected by off-the-energy-shell effects

## Technical issue II - Recoil corrections at N<sup>3</sup>LO

$$\mathbf{j}^{\text{N}^3\text{LO}} =$$

► Reducible contributions


$$\begin{aligned} \mathbf{j}^{\text{red}} &\sim \int v^\pi(\mathbf{q}_2) \frac{1}{E_i - E_f} \mathbf{j}^{\text{NLO}}(\mathbf{q}_1) \\ &- \int 2 \frac{\omega_1 + \omega_2}{\omega_1 \omega_2} V_{\pi NN}(2, \mathbf{q}_2) V_{\pi NN}(2, \mathbf{q}_1) V_{\pi NN}(1, \mathbf{q}_2) V_{\gamma\pi NN}(1, \mathbf{q}_1) \end{aligned}$$

► Irreducible contributions

$$\begin{aligned} \mathbf{j}^{\text{irr}} &= \int 2 \frac{\omega_1 + \omega_2}{\omega_1 \omega_2} V_{\pi NN}(2, \mathbf{q}_2) V_{\pi NN}(2, \mathbf{q}_1) V_{\pi NN}(1, \mathbf{q}_2) V_{\gamma\pi NN}(1, \mathbf{q}_1) \\ &- \int 2 \frac{\omega_1^2 + \omega_2^2 + \omega_1 \omega_2}{\omega_1 \omega_2 (\omega_1 + \omega_2)} [V_{\pi NN}(2, \mathbf{q}_2), V_{\pi NN}(2, \mathbf{q}_1)]_- V_{\pi NN}(1, \mathbf{q}_2) V_{\gamma\pi NN}(1, \mathbf{q}_1) \end{aligned}$$

► Observed partial cancellations at N<sup>3</sup>LO between recoil corrections to reducible diagrams and irreducible contributions

## The box diagram: an example at N<sup>3</sup>LO




$$\begin{aligned}
 \text{direct} &= f_d(\omega_1, \omega_2) V_a V_b V_c V_d \\
 \text{crossed} &= f_c(\omega_1, \omega_2) V_b V_a V_c V_d \qquad V_b V_a = V_a V_b - [V_a, V_b]_-
 \end{aligned}$$

$$\begin{aligned}
 \text{irreducible} &= [f_d(\omega_1, \omega_2) + f_c(\omega_1, \omega_2)] V_a V_b V_c V_d \\
 &- f_c(\omega_1, \omega_2) [V_a, V_b]_- V_c V_d
 \end{aligned}$$

## Transition amplitude in time-ordered perturbation theory

$$\begin{aligned}
 T_{fi} = \langle f | T | i \rangle &= \langle f | H_1 \sum_{n=1}^{\infty} \left( \frac{1}{E_i - H_0 + i\eta} H_1 \right)^{n-1} | i \rangle \\
 &= \langle f | H_1 | i \rangle + \sum_{|I\rangle} \langle f | H_1 | I \rangle \frac{1}{E_i - E_I} \langle I | H_1 | i \rangle + \dots
 \end{aligned}$$

- ▶ A contribution with N interaction vertices and L loops scales as

$$\underbrace{e \left( \prod_{i=1}^N Q^{\alpha_i - \beta_i / 2} \right)}_{H_1 \text{ scaling}} \times \underbrace{Q^{-(N - N_K - 1)} Q^{-2N_K}}_{\text{denominators}} \times \underbrace{Q^{3L}}_{\text{loop integration}}$$


$\alpha_i$  = number of derivatives in  $H_1$  and  $\beta_i$  = number of  $\pi$ 's at each vertex

$N_K$  = number of pure nucleonic intermediate states


- ▶  $(N - N_K - 1)$  energy denominators expanded in powers of  $(E_i - E_N)/\omega_\pi \sim Q$

$$\frac{1}{E_i - E_I} |I\rangle = \frac{1}{E_i - E_N - \omega_\pi} |I\rangle \sim - \left[ \underbrace{\frac{1}{\omega_\pi}}_{Q^{-1}} + \underbrace{\frac{E_i - E_N}{\omega_\pi^2}}_{Q^0} + \underbrace{\frac{(E_i - E_N)^2}{\omega_\pi^3}}_{Q^1} + \dots \right] |I\rangle$$

- ▶ Due to the chiral expansion, the transition amplitude  $T_{fi}$  can be expanded as

$$T_{fi} = T^{\text{LO}} + T^{\text{NLO}} + T^{\text{N2LO}} + \dots \quad \text{and} \quad T^{\text{NnLO}} \sim (Q/\Lambda_\chi)^n T^{\text{LO}}$$

## EM charge up to $n = 0$ (or up to N3LO)



▶  $n = -3$

$$\rho^{(-3)}(\mathbf{q}) = e(2\pi)^3 \delta(\mathbf{p}_1 + \mathbf{q} - \mathbf{p}'_1) (1 + \tau_{1,z})/2 + 1 \Rightarrow 2$$

▶  $n = -1$ :

$$(Q/m_N)^2 \text{ relativistic correction to } \rho^{(-3)}$$

▶  $n = 0$ :

i) 'static' tree-level current (originates from a  $\gamma\pi N$  vertex of order  $eQ$ )

ii) 'non-static' OPE charge operators,  $\rho_\pi^{(0)}(\mathbf{v})$  depends on  $v_\pi^{(2)}(\mathbf{v})$


▶  $\rho_\pi^{(0)}(\mathbf{v})$ 's are unitarily equivalent

$$\rho_\pi^{(0)}(\mathbf{v}) = \rho_\pi^{(0)}(\mathbf{v} = 0) + [\rho^{(-3)}, iU^{(0)}(\mathbf{v})]$$

▶ No unknown LECs up to this order ( $g_A, F_\pi$ )


EM charge @  $n = 1$  (or N4LO)

1.



- ▶ (a), (f), (d), and (i) vanish
- ▶ Divergencies associated with (b) + (g), (c) + (h), and (e) + (j) cancel out—as they must since there are no counter-terms at N4LO
- ▶  $\rho_{\text{h}}^{(1)}(\mathbf{v})$  depends on the parametrization adopted for  $v_{\pi}^{(2)}(\mathbf{v})$  and  $v_{2\pi}^{(3)}(\mathbf{v})$
- ▶  $\rho_{\text{h}}^{(1)}(\mathbf{v})$ 's are unitarily equivalent

$$\rho_{\text{h}}^{(1)}(\mathbf{v}) = \rho_{\text{h}}^{(1)}(\mathbf{v} = 0) + [\rho^{(-3)}, iU^{(1)}(\mathbf{v})]$$



- Charge operators ( $\mathbf{v}$ -dependent included) up to  $n = 1$  satisfy the condition

$$\rho^{(n>-3)}(\mathbf{q} = 0) = 0$$

which follows from charge conservation

$$\rho(\mathbf{q} = 0) = \int d\mathbf{x} \rho(\mathbf{x}) = e \frac{(1 + \tau_{1,z})}{2} + 1 \Leftrightarrow 2 = \rho^{(-3)}(\mathbf{q} = 0)$$

- $\rho^{(1)}$  does not depend on unknown LECs and it is purely isovector