


Eigenvector continuation and new results in nuclear lattice simulations

Dean Lee

Facility for Rare Isotope Beams
Department of Physics and Astronomy
Michigan State University
Nuclear Lattice EFT Collaboration

Progress in *Ab Initio* Techniques in Nuclear Physics
TRIUMF, February 28, 2018



Outline

Lattice effective field theory


Pinhole algorithm

Pinhole trace algorithm


Eigenvector continuation

Summary and outlook

Lattice chiral effective field theory



Review: D.L, Prog. Part. Nucl. Phys. 63 117-154 (2009)
TALENT summer school lectures: qmc2016.wordpress.ncsu.edu



Challenge

In order to compute thermodynamic properties of finite nuclei, nuclear matter, and neutron matter, we need to compute the partition function

$$\text{Tr exp}(-\beta H)$$

The standard method for computing the partition function involves calculating determinants of matrices of size $4V \times 4V$, where V is the number of lattice points filling the spatial volume. Since V is usually several hundred or several thousand, these calculations are very expensive.

Pinhole algorithm

Consider the density operator for nucleon with spin i and isospin j


$$\rho_{i,j}(\mathbf{n}) = a_{i,j}^\dagger(\mathbf{n})a_{i,j}(\mathbf{n})$$


We construct the normal-ordered A -body density operator

$$\rho_{i_1,j_1,\dots,i_A,j_A}(\mathbf{n}_1,\dots,\mathbf{n}_A) = : \rho_{i_1,j_1}(\mathbf{n}_1) \cdots \rho_{i_A,j_A}(\mathbf{n}_A) :$$

In the simulations we do Monte Carlo sampling of the amplitude

$$A_{i_1,j_1,\dots,i_A,j_A}(\mathbf{n}_1,\dots,\mathbf{n}_A,t) = \langle \Psi_I | e^{-Ht/2} \rho_{i_1,j_1,\dots,i_A,j_A}(\mathbf{n}_1,\dots,\mathbf{n}_A) e^{-Ht/2} | \Psi_I \rangle$$





Elhatisari, Epelbaum, Krebs, Lähde, D.L., Li, Lu, Meißner, Rupak, PRL 119, 222505 (2017)

Pinhole trace algorithm


We have developed an alternative method using pinholes that calculates determinants of matrices of size $A \times A$, where A is the number of nucleons. The method does not suffer from severe sign oscillations.

We compute the quantum mechanical trace over A -nucleon states by summing over pinholes (position eigenstates) for the initial and final states

$$\begin{aligned} & \text{Tr } O \\ &= \frac{1}{A!} \sum_{i_1 \cdots i_A, j_1 \cdots j_A, \mathbf{n}_1 \cdots \mathbf{n}_A} \langle 0 | a_{i_A, j_A}(\mathbf{n}_A) \cdots a_{i_1, j_1}(\mathbf{n}_1) O a_{i_1, j_1}^\dagger(\mathbf{n}_1) \cdots a_{i_A, j_A}^\dagger(\mathbf{n}_A) | 0 \rangle \end{aligned}$$

This can be used to calculate the partition function in the canonical ensemble.

Metropolis updates of pinholes






Figure courtesy of Bingnan Lu




Figure courtesy of Bingnan Lu

Challenge

A common challenge faced in many fields of quantum physics is finding the extremal eigenvalues and eigenvectors of a Hamiltonian matrix too large to store in computer memory.

There are numerous efficient methods developed for this task. All existing methods either use Monte Carlo simulations, diagrammatic expansions, variational methods, or some combination.

The problem is that they generally fail when some control parameter in the Hamiltonian matrix exceeds some threshold value.

Eigenvector continuation

We demonstrate that when a control parameter in the Hamiltonian matrix is varied smoothly, the extremal eigenvectors do not explore the large dimensionality of the linear space. Instead they trace out trajectories with significant displacements in only a small number of linearly-independent directions.

We prove this empirical observation using analytic function theory and the principles of analytic continuation.

Since the eigenvector trajectory is a low-dimensional manifold embedded in a very large space, we can find the desired eigenvector using methods similar to image recognition in machine learning.

D. Frame, R. He, I. Ipsen, Da. Lee, De. Lee, E. Rrapaj, arXiv:1711.07090

Consider a one-parameter family of Hamiltonian matrices of the form

$$H(c) = H_0 + cH_1$$

where H_0 and H_1 are Hermitian. Let the eigenvalues and eigenvectors be

$$H(c)|\psi_j(c)\rangle = E_j(c)|\psi_j(c)\rangle$$


We can perform series expansions around the point $c = 0$.

$$E_j(c) = \sum_{n=0}^{\infty} E_j^{(n)}(0)c^n/n!$$
$$|\psi_j(c)\rangle = \sum_{n=0}^{\infty} |\psi_j^{(n)}(0)\rangle c^n/n!$$

This is the strategy of perturbation theory. We can compute each term in the series when the eigenvalues and eigenvectors of H_0 are known or computable.

Perturbation theory

$$|\psi_j(c)\rangle = \sum_{n=0}^{\infty} |\psi_j^{(n)}(0)\rangle c^n / n!$$



Bose-Hubbard model


In order to illuminate our discussion with a concrete example, we consider a quantum Hamiltonian known as the Bose-Hubbard model in three dimensions. It describes a system of identical bosons on a three-dimensional cubic lattice.

$$H = -t \sum_{\langle \mathbf{n}', \mathbf{n} \rangle} a^\dagger(\mathbf{n}') a(\mathbf{n}) + \frac{U}{2} \sum_{\mathbf{n}} \rho(\mathbf{n}) [\rho(\mathbf{n}) - \mathbf{1}] - \mu \sum_{\mathbf{n}} \rho(\mathbf{n})$$
$$\rho(\mathbf{n}) = a^\dagger(\mathbf{n}) a(\mathbf{n})$$


The parameter t controls the hopping the bosons on the lattice, and U is the single-site pairwise interaction. We set the chemical potential to be

$$\mu = -6t$$


Perturbation theory fails at strong attractive coupling




Restrict the linear space to the span of three vectors




$$|\psi_j(c)\rangle = \sum_{n=0}^{\infty} |\psi_j^{(n)}(w)\rangle (c-w)^n / n!$$






$$|\psi_j(c)\rangle = \sum_{n=0}^{\infty} |\psi_j^{(n)}(w)\rangle (c-w)^n / n!$$

$$|\psi_j^{(n)}(w)\rangle = \sum_{m=0}^{\infty} |\psi_j^{(n+m)}(0)\rangle w^m / m!$$



$$|\psi_j(c)\rangle = \sum_{n=0}^{\infty} |\psi_j^{(n)}(w)\rangle (c-w)^n / n!$$

$$|\psi_j^{(n)}(w)\rangle = \sum_{m=0}^{\infty} |\psi_j^{(n+m)}(0)\rangle w^m / m!$$



$$|\psi_j(c)\rangle = \sum_{n=0}^{\infty} |\psi_j^{(n)}(0)\rangle c^n / n!$$

$$|\psi_j(c)\rangle = \lim_{N, M \rightarrow \infty} \sum_{n=0}^N \sum_{m=0}^M |\psi_j^{(n+m)}(0)\rangle w^m (c-w)^n / (m! n!)$$

The eigenvector can be well approximated as a linear combination of a few vectors, using either the original series expansion


$$|\psi_j(c)\rangle = \sum_{n=0}^{\infty} |\psi_j^{(n)}(0)\rangle c^n / n!$$

or the rearranged multi-series expansion we obtained through analytic continuation


$$|\psi_j(c)\rangle = \lim_{N, M \rightarrow \infty} \sum_{n=0}^N \sum_{m=0}^M |\psi_j^{(n+m)}(0)\rangle w^m (c-w)^n / (m! n!)$$

As c is varied the eigenvector does not explore the large dimensionality of the linear space, but is instead well approximated by a low-dimension manifold.

We can “learn” the eigenvector trajectory in one region and perform eigenvector continuation to another region




Applying eigenvector continuation to more than one eigenvector at a time accelerates convergence near avoided level crossings.



Application: Neutron matter simulations


We consider lattice effective field theory simulations of the neutron matter at the leading order.




As a challenge to the eigenvector continuation technique, we use a lattice action for one-pion exchange that causes severe Monte Carlo sign oscillations.

D.L., in “An Advanced Course in Computational Nuclear Physics”, Hjorth-Jensen, Lombardo, van Kolck, Eds., Lecture Notes in Physics, Volume 936 [arXiv:1609.00421]


Direct calculation of six neutrons ($L = 8$ fm)




Direct calculation of fourteen neutrons ($L = 8$ fm)



Eigenvector continuation for six neutrons ($L = 8$ fm)



Eigenvector continuation for fourteen neutrons ($L = 8$ fm)



g_A^2 values	E_0 for six neutrons (MeV)	E_0 for fourteen neutrons (MeV)
c_1	13.8(1)	48.9(4)
c_2	13.6(2)	48.4(5)
c_3	13.6(2)	48.9(6)
c_2, c_3	13.6(2)	48.1(6)
c_3, c_1	13.6(2)	48.9(6)
c_1, c_2	13.6(2)	48.0(6)
c_1, c_2, c_3	13.6(2)	48.0(6)
direct calculation	$12\begin{pmatrix} +3 \\ -4 \end{pmatrix}$	$42\begin{pmatrix} +7 \\ -15 \end{pmatrix}$

Summary and Outlook

These are exciting times for the *ab initio* nuclear theory community. In lattice EFT, we have new projects in motion which are pushing the current frontiers.

Currently working to improve our understanding of the detailed connection between bare nuclear forces and nuclear structure for light and medium-mass nuclei.

Applying the adiabatic projection method to low-energy nucleon-nucleus and alpha-nucleus scattering and reactions.

Using the pinhole algorithm to study the detailed structure of nuclei and pinhole trace algorithm for thermodynamics of finite nuclei, nuclear matter, and neutron matter.

Implementing eigenvector continuation to treat all higher-order interactions in chiral effective field theory.