

Towards Accurate Calculations of Medium-Mass Nuclei


Robert Roth



TECHNISCHE
UNIVERSITÄT
DARMSTADT

10th TRIUMF Ab Initio Workshop

we have a powerful and versatile many-body toolbox at our disposal...



...many-body uncertainties are under control

...chiral inputs for interactions and operators limit accuracy

Ab Initio Toolbox

light nuclei

NCSM

- solution of matrix eigenvalue problem in truncated many-body model space
- uncertainties: N_{\max} convergence

In-Medium NCSM

- NCSM calculation with an in-medium SRG decoupled many-body Hamiltonian
- uncertainties: NO2B, $E_{3\max}$, e_{\max} , N_{\max} convergence

In-Medium SRG

- decoupling ground-state from excitations through unitary transformation via flow equations
- uncertainties: NO2B initial and flow equation, $E_{3\max}$, e_{\max}

Coupled Cluster

- decoupling ground-state from excitations through similarity transformation via nonlinear amplitude equations
- uncertainties: triples correction, NO2B initial, $E_{3\max}$, e_{\max}

heavy nuclei

MBPT, SCGF, ...

Ab Initio Toolbox - Add-Ons

Free-Space SRG

- momentum-space decoupling of initial Hamiltonian through unitary transformation
- consistent transformation of $NN+3N$ interaction and all relevant operators
- uncertainties: induced many-body contributions

Basis Optimization

- Hartree-Fock basis
- natural orbitals from perturbatively improved density matrix
- uncertainties: $NO2B$, E_{3max} , e_{max}

multiple sources of uncertainties...

...quantification not trivial

Third Generation Chiral Interactions

A Brief History... Incomplete and Totally Biased

1st Generation

- 2007:** first ab initio calculation of mid-p-shell nuclei with local chiral 3N interaction: $N3LO_{EM} + N2LO_{L,500}$ *PRL 99, 042501 (2007)*
- 2012:** SRG transformed NN+3N interactions and reduced 3N cutoffs for oxygen & calcium isotopes *PRL 109, 052501 (2012)*
- 2014:** overbinding beyond oxygen and catastrophic radii *PLB 736, 119 (2014)*

2nd Generation

- 2015:** combined fit of few and many-body observables to improve radii, sacrificing phase-shifts: $N2LO_{SAT}$ *PRC 91, 051301(R) (2015)*
- 2016:** magic interactions constructed from a SRG evolved NN interaction plus bare 3N parametrization *PRC 83, 031301(R) (2011)*
PRC 93, 011302 (2016)

3rd Generation

- 2016:** systematic order-by-order calculations up to $N3LO$ of neutron and nuclear matter *PRC 94, 054307 (2016)*
PRL 122, 042501 (2019)
- 2019:** systematic order-by-order calculations up to $N3LO$ in light and medium-mass nuclei... *PRC 96, 024004 (2017)*
arXiv:1911.04955 (2019)


Nuclear Interactions from Chiral EFT

NN


LO




NLO




N2LO



N3LO



N4LO



NN @ LO, NLO,
 N2LO, N3LO
 Entem, Machleidt & Nosyk,
 non-local regulator,
 cutoff 450, 500, 550 MeV

 3N @ N2LO, N3LO
 non-local regulator,
 cutoff 450, 500, 550 MeV

3rd Generation

- NN+3N interactions with consistent chiral orders and cutoffs
- vary the chiral order and the cutoff systematically to assess uncertainties
- explore dependence on regulator scheme (local, non-local, semi-local)

LENPIC:
 analogous family
 with semi-local
 regulators

■ start from chiral NN interaction by Entem, Machleidt & Nosyk

PRC 96, 024004 (2017)

- LO to N3LO
- non-local regulator
- cutoff 450, 500, 550 MeV
- accurate reproduction of NN scattering data up to ~ 300 MeV

■ supplement non-local 3N interaction at N2LO and N3LO


- N2LO or N3LO, consistent with NN interaction
- non-local regulator, as in NN interaction
- cutoff 450, 500, 550 MeV, consistent with NN interaction

■ fix c_E in few-body sector, keep c_D as a parameter

- c_E fit to triton binding energy
- alternative: c_E from combined fit to ${}^3\text{H}$, ${}^4\text{He}$ energy and ${}^4\text{He}$ radius

^4He Ground State: c_D Scan


Hüther et al.; arXiv:1911.04955



- Jacobi-NCSM calculations for ^3H and ^4He with bare interaction
- scanning c_D over large range, c_E always fit to ^3H binding energy

^4He Ground State: c_D Scan


Hüther et al.; arXiv:1911.04955




- Jacobi-NCSM calculations for ^3H and ^4He with bare interaction
- scanning c_D over large range, c_E always fit to ^3H binding energy

Medium-Mass Nuclei: c_D Scan

Hüther et al.; arXiv:1911.04955




Non-Local vs. Local 3N Regulator



Uncertainties

Interaction Uncertainties

Hüther et al.; arXiv:1911.04955



- quantify uncertainties from order-by-order systematics
- simplified protocol based on expansion parameter $Q=q/\Lambda_B$

$$\delta X_{N3LO} = \max($$

$$Q |X_{N3LO} - X_{N2LO}|,$$

$$Q^2 |X_{N2LO} - X_{NLO}|,$$

$$Q^4 |X_{NLO} - X_{LO}|,$$

$$Q^5 |X_{LO}|)$$


PRC 98, 014002 (2018)
PRC 93, 044002 (2016)

IM-SRG(M2)
natural orbitals
 $\hbar\Omega=20$ MeV
 $e_{\max}=12, E_{3\max}=16$
 $\alpha=0.04$ fm⁴




Many-Body Uncertainties I

Hüther et al.; arXiv:1911.04955




Many-Body Uncertainties II

Hüther et al.; arXiv:1911.04955



Medium-Mass Nuclei

Hüther et al.; arXiv:1911.04955




IM-SRG(M2), natural orbitals, $\hbar\Omega=20$ MeV, $\alpha=0.04$ fm⁴, $e_{\text{max}}=12$, $E_{3\text{max}}=16$

error bands show interaction uncertainties

Medium-Mass Nuclei


Hüther et al.; arXiv:1911.04955



IM-SRG(M2), natural orbitals, $\hbar\Omega = 20$ MeV, $\alpha = 0.04$ fm⁴, $e_{\text{max}} = 12$, $E_{3\text{max}} = 16$

Oxygen Isotopic Chain


Hüther et al.; arXiv:1911.04955




IM-NCSM, natural orbitals, $\hbar\Omega = 20$ MeV, $\alpha = 0.04$ fm⁴, $e_{\max} = 12$, $E_{3\max} = 14$, $N_{\text{ref}} = 2$

p-Shell Spectra


⁷Li




⁸Be




⁹Be



¹⁰B



¹²C



NCSM/IM-NCSM, $\Lambda=500$ MeV, $\hbar\Omega=20$ MeV
error bands show interaction uncertainties

see poster by
Tobias Mongelli


Uncertain Uncertainties

-or-

The “Expert Assessment”

2%... really?

Many-Body Uncertainties



based on energies computed with a specific class of SRG-transformed Hamiltonians

does this hold for bare calculations, other interactions or other observables ???



$N3LO_{EM} + N2LO_{L,400}$
 $\alpha = 0.08 \text{ fm}^4$
 natural orbitals / HF
 $\hbar\Omega = 20 \text{ MeV}$
 $e_{\max} = 12$
 $N_{\max} = 0$ reference

A Simple Exercise

revisit
many-body uncertainties
through a very simple
exercise...

- compare a few standard medium-mass calculations for ground-state energies and radii and compare results with...
 - **bare and SRG-transformed Hamiltonians** starting from
 $N2LO_{SAT}$ and $N3LO_{EMN,500} + N3LO_{NL,500}$
 - different **medium-mass methods**
IM-SRG(M2), CCSD, CR-CC(2,3), Λ -CCSD(T)
 - different **single-particle bases**
Hartree-Fock, natural orbitals, more creative


^{16}O with $\text{N}^2\text{LO}_{\text{SAT}}$



- ◆ standard HF basis for actual Hamiltonian
- standard NAT basis for actual Hamiltonian
- ◆ HF basis for SRG-evolved Hamiltonian with $\alpha=0.08 \text{ fm}^4$
- NAT basis for SRG-evolved Hamiltonian with $\alpha=0.08 \text{ fm}^4$

basis variation provides simple tool to probe uncertainties due to reference state, NO^2B , $E_{3\text{max}}, \dots$

^{16}O with $\text{N}3\text{LO}_{\text{EMN},500} + \text{N}3\text{LO}_{\text{NL},500}$



- ◆ standard HF basis for actual Hamiltonian
- standard NAT basis for actual Hamiltonian
- ◆ HF basis for SRG-evolved Hamiltonian with $\alpha=0.08 \text{ fm}^4$
- NAT basis for SRG-evolved Hamiltonian with $\alpha=0.08 \text{ fm}^4$

Summary

some solutions

...new family of NN+3N interactions up to N3LO with non-local regulators for different cutoffs

...quantification of interaction and many-body uncertainties

...excellent description of ground-state energies, radii, spectra

more questions

...why the drastic differences between local and non-local 3N regulators ?

...why the different optimal c_D values in few-body/many-body/matter ?

...are we sometimes overly optimistic with our many-body uncertainties ?

Epilogue

■ thanks to my group and my collaborators

- S. Alexa, **T. Hüther**, M. Knöll, L. Mertes, T. Mongelli, J. Müller, K. Vobig, T. Wolfgruber & **K. Hebler**, A. Tichai
Technische Universität Darmstadt
- T. Duguet
CEA Saclay
- P. Navrátil
TRIUMF, Vancouver
- H. Hergert, R. Wirth
NSCL / Michigan State University
- J. Vary, P. Maris
Iowa State University
- R. Machleidt
University of Idaho
- E. Epelbaum, H. Krebs, P. Reinhard & the LENPIC Collaboration
Universität Bochum, ...



Deutsche
Forschungsgemeinschaft

DFG



Exzellente Forschung für
Hessens Zukunft

